OREGON IMAGING CENTERS

Today's Date: _____

CARDIOVASCULAR CT ORDER FORM

General Scheduling: (541) 334-7555
Toll free: (888) 968-7608 ◆ **Fax Orders: (541) 334-7564**
Online Orders: *https://scheduling.oregonimaging.com*

All Coronary CT exams are performed at our University District location:
1200 Hilyard St., Suite 330
Eugene, OR 97401

PATIENT LAST NAME **(REQUIRED)**	FIRST	M	DATE OF BIRTH **(REQUIRED)**	PATIENT'S PHONE
ORDERING CLINICIAN **(REQUIRED)** OFFICE LOCATION (if multiple)		CLINICIAN SIGNATURE **(REQUIRED)**		PROVIDER PHONE AND FAX
SEND ADDITIONAL COPIES OF REPORT TO		INSURANCE AND MEMBER ID#		PA#

Reason for Exam/ ICD10 Code: _____

☐ I20.8 Other forms angina pectoris ☐ I25.10 CAD w/o angina ☐ R94.3 Abnormal result CV function study

☐ I11.9 Hypertensive Heart disease ☐ R06.02 Shortness of breath ☐ I42.9 Cardiomyopathy, unspecified

CORONARY/CARDIAC CT

☐ Coronary CTA with rate controlling medication (75574)
- ☐ FFRct* (75580)
- ☐ Plaque Analysis* (75577)

*FFRct and Plaque Analysis only performed when indicated in accordance with current guidelines

☐ Coronary Artery Calcium Score (75571)

Coronary CTA Exam Prep Instructions:

- No food for 4 hours prior to the exam
- Drink plenty of water or clear fluids without caffeine up until time of exam (patient should be well hydrated for renal protection, for ease of establishing venous access, and to avoid post-procedure hypotension).
- No caffeine products for 12 hours prior to exam because they might hinder efforts to reduce the heart rate before scanning. This includes coffee, tea, energy drinks, energy pills, diet pills, and most soda.
- Avoid exercise and nicotine for 8 hours before scan.
- Avoid medications for Erectile Disfunction or Female Sexual Arousal Disorder such as Viagra or Cialis for 72 hours before exam.
- Take all other regular medications the day of the exam, especially blood pressure medicine.

Medication Prep for Coronary CTA Studies:

In order to decrease cardiac motion and improve diagnostic accuracy, a heart rate of 60 bpm or less is optimal at the time of image acquisition.

Suggested Premedication Protocol:

- Medication for resting heart rate 60-69 bpm; 50mg Metoprolol Tartrate by mouth one hour before scheduled scan time.
- Medication for resting heart rate ≥70 bpm; 100 mg Metoprolol Tartrate by mouth one hour before scheduled scan time.
- **Pre-scan medications should be prescribed by referring clinician**. Contraindications to Metoprolol include sinus bradycardia, severe aortic stenosis, allergy to beta blockers, COPD/Asthma requiring use of inhaler, heart block, uncontrolled heart failure or blood pressure less than 100/60 mmHg.
- **Irregular heart rate is contraindicated**.